

The crossing numbers of  $K_{n,n} - nK_2$ ,  $K_n \times P_2$ ,  $K_n \times P_3$  and

$$K_n \times C_4^*$$

Yuansheng Yang<sup>†</sup>, Baigong Zheng, Xiaohui Lin, Xirong Xu

School of Computer Science and Technology

Dalian University of Technology, Dalian, 116024, P.R. China

### Abstract

The *crossing number* of a graph  $G$  is the minimum number of pairwise intersections of edges among all drawings of  $G$ . In this paper, we study the crossing number of  $K_{n,n} - nK_2$ ,  $K_n \times P_2$ ,  $K_n \times P_3$  and  $K_n \times C_4$ .

**Keywords:** *Crossing number; Drawing; Complete bipartite graphs; Kronecker product*

## 1 Introduction and preliminaries

Let  $G$  be a graph,  $V(G)$  the vertex set and  $E(G)$  the edge set. The *crossing number* of  $G$ , denoted by  $cr(G)$ , is the smallest number of pairwise crossings of edges among all drawings of  $G$  in the plane. We use  $D$  to denote a drawing of a graph  $G$  and  $\nu(D)$  the number of crossings in  $D$ . It is clear that  $cr(G) \leq \nu(D)$ .

Let  $E_1$  and  $E_2$  be two disjoint subsets of an edge set  $E$ . The number of the crossings formed by an edge in  $E_1$  and another edge in  $E_2$  is denoted by  $\nu_D(E_1, E_2)$  in a drawing  $D$ . The number of the crossings that involve a pair of edges in  $E_1$  is denoted by  $\nu_D(E_1)$ . Then  $\nu(D) = \nu_D(E(G))$  and  $\nu_D(E_1 \cup E_2) = \nu_D(E_1) + \nu_D(E_2) + \nu_D(E_1, E_2)$ .

The *Kronecker product*  $G \times H$  of graphs  $G$  and  $H$  has vertex set  $V(G \times H) = V(G) \times V(H)$  and edge set  $E(G \times H) = \{(a, x), (b, y)\} : \{a, b\} \in E(G) \text{ and } \{x, y\} \in E(H)\}$ .

---

\*The research is supported by NSFC of China (No.60973014, 61170303)

<sup>†</sup>corresponding author's email : yangys@dlut.edu.cn

(The product is also known as direct product, cardinal product, cross product and graph conjunction.)

Computing the crossing number of graphs is a complicated yet classical problem. It is proved that the problem is NP-complete by Garey and Johnson [7].

Zarankiewicz [14] gave a drawing of  $K_{m,n}$ , which demonstrated that

$$cr(K_{m,n}) \leq Z(m, n) = \lfloor \frac{m}{2} \rfloor \lfloor \frac{m-1}{2} \rfloor \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor.$$

The equality holds for  $\min(m, n) \leq 6$  [11] and  $7 \leq n \leq 8, 7 \leq m \leq 10$  [13].

Gue [8] and Blažek and Koman [4] gave a drawing of  $K_n$  in a cylinder (homeomorphic to a plane, a sphere), which demonstrated that

$$cr(K_n) \leq Z(n) = \frac{1}{4} \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor.$$

The equality holds for  $n \leq 12$  [2].

De Klerk, Pasechnik and Schrijver [5] proved  $cr(K_{m,n}) \geq 0.8594Z(m, n)$  and  $cr(K_n) \geq 0.8594Z(n)$ .

In literature, the Cartesian product has been paid more attention [1, 15–17], while Kronecker product has fewer results on the crossing number [9].

In this paper, we study the crossing numbers of  $K_{n,n} - nK_2$  and the Kronecker product  $K_n \times P_2, K_n \times P_3, K_n \times C_4$ . In Section 2, we give upper bounds of  $cr(K_{n,n} - nK_2)$  and  $cr(K_n \times P_2)$  by constructing a drawing of  $K_{n,n} - nK_2$  in a cylinder. In Section 3, we give upper bound of  $cr(K_n \times P_3)$  by constructing a drawing of  $K_n \times P_3$  based on the drawing of  $K_{n,n} - nK_2$ . In Section 4, we give upper bound of  $cr(K_n \times C_4)$  by constructing a drawing of  $K_n \times C_4$  based on the drawing of  $K_{n,n} - nK_2$ , too. In Section 5, we give lower bounds of  $cr(K_{n,n} - nK_2), cr(K_n \times P_2), cr(K_n \times P_3)$  and  $cr(K_n \times C_4)$ .

## 2 Upper bounds of $cr(K_{n,n} - nK_2)$ and $cr(K_n \times P_2)$

Let

$$\begin{aligned} V(K_{n,n} - nK_2) &= \{a_i, b_i \mid 0 \leq i \leq n-1\}, \\ E(K_{n,n} - nK_2) &= \{(a_i, b_j) \mid 0 \leq i \neq j \leq n-1\}. \end{aligned}$$

In Figure 2.1, we exhibit drawings  $D_n$  of  $K_{n,n} - nK_2$  in a cylinder for  $n \leq 13$ . A




Figure 2.1: Drawings  $D_n$  for  $n \leq 13$

cylinder can be ‘assembled’ from a polygon by identifying one pair of opposite sides of a rectangle [3].

From drawings in Figure 2.1, we have

**Lemma 2.1.** *For  $n \leq 4$ ,  $cr(K_{n,n} - nK_2) = 0$ .*

For even  $n \geq 6$ , let

$$\begin{aligned}
 X &= \{a_{2i}, b_{2i+1} \mid 0 \leq i \leq n/2 - 1\}, \\
 Y &= \{a_{2i+1}, b_{2i} \mid 0 \leq i \leq n/2 - 1\}, \\
 E_X &= \{uv \mid u, v \in X\}, \\
 E_{XY} &= \{uv \mid u \in X \text{ and } v \in Y\}, \\
 E_Y &= \{uv \mid u, v \in Y\}.
 \end{aligned}$$

Then

$$\begin{aligned}
 V(K_{n,n} - nK_2) &= X \cup Y, \quad X \cap Y = \emptyset. \\
 E(K_{n,n} - nK_2) &= E_X \cup E_{XY} \cup E_Y.
 \end{aligned}$$

In drawings  $D_n$  with even  $n \geq 6$ , vertices  $a_{2\lceil \frac{n}{4} \rceil - 1}, b_{2\lceil \frac{n}{4} \rceil}, \dots, a_{2\lceil \frac{n}{4} \rceil - 3}, b_{2\lceil \frac{n}{4} \rceil - 2}$  ( $a_0, b_1, \dots, a_{n-2}, b_{n-1}$ ) of  $X$  ( $Y$ ) are placed equidistantly on the perimeter of the top (bottom) disk, edges of  $E_X$  ( $E_Y$ ) are drawn on the top (bottom) disk, and edges of  $E_{XY}$  are drawn by shortest helical curves on the cylinder. It follows  $\nu_{D_n}(E_X, E_{XY}) = \nu_{D_n}(E_Y, E_{XY}) = \nu_{D_n}(E_X, E_Y) = 0$  and  $\nu(D_n) = \nu_{D_n}(E_X) + \nu_{D_n}(E_Y) + \nu_{D_n}(E_{XY}) = 2\nu_{D_n}(E_X) + \nu_{D_n}(E_{XY})$ .

For even  $n \geq 6$  and  $n \bmod 4=2$ ,

$$\begin{aligned}
\nu_{D_n}(E_X) &= \frac{1}{2} \left( n \sum_{d=1}^{\frac{n-6}{4}} 2d \left( \frac{n}{2} - 1 - d \right) + \frac{n}{2} 2 \left( \frac{n-2}{4} \right)^2 \right) \\
&= \frac{n(n-2)}{2} \frac{\frac{n-6}{4} \frac{n-2}{4}}{2} - n \frac{\frac{n-6}{4} \frac{n-2}{4} \frac{n-4}{2}}{6} + \frac{n(n-2)^2}{32} \\
&= \frac{n(n-2)(3n^2-24n+36-n^2+10n-24+6n-12)}{192} \\
&= \frac{n^2(n-2)(n-4)}{96}.
\end{aligned}$$

For even  $n \geq 8$  and  $n \bmod 4=0$ ,

$$\begin{aligned}
\nu_{D_n}(E_X) &= \frac{n}{2} \sum_{d=1}^{\frac{n-4}{4}} 2d \left( \frac{n}{2} - 1 - d \right) \\
&= \frac{n(n-2)}{2} \frac{\frac{n}{4} \frac{n-4}{4}}{2} - n \frac{\frac{n-4}{4} \frac{n}{4} \frac{n-2}{2}}{6} \\
&= \frac{n^2(n-2)(n-4)}{64} - \frac{n^2(n-2)(n-4)}{192} \\
&= \frac{n^2(n-2)(n-4)}{96}.
\end{aligned}$$

For even  $n \geq 6$ ,

$$\begin{aligned}
\nu_{D_n}(E_{XY}) &= n \sum_{d=1}^{\frac{n-4}{2}} \sum_{i=1}^d (2i-1) \\
&= n \sum_{d=1}^{\frac{n-4}{2}} d^2 \\
&= n \frac{\frac{n-4}{2} \frac{n-2}{2} (n-3)}{6} \\
&= \frac{n(n-2)(n-3)(n-4)}{24}.
\end{aligned}$$

Hence, for even  $n \geq 6$ ,

$$\begin{aligned}
\nu(D_n) &= 2\nu_{D_n}(E_X) + \nu_{D_n}(E_{XY}) \\
&= \frac{n^2(n-2)(n-4)}{48} + \frac{n(n-2)(n-3)(n-4)}{24} \\
&= \frac{n(n-2)^2(n-4)}{16} \\
&= \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor.
\end{aligned}$$

So we have the following Lemma 2.2:

**Lemma 2.2.** For even  $n \geq 6$ ,  $\nu(D_n) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ .

For odd  $n \geq 5$ , let

$$\begin{aligned}
X &= \{a_{2i}, b_{2i+1} \mid 0 \leq i < (n-1)/2\}, \\
Y &= \{a_{2i+1}, b_{2i} \mid 0 \leq i < (n-1)/2\}, \\
Z &= \{a_{n-1}, b_{n-1}\}, \\
E_X &= \{uv \mid u, v \in X\}, \\
E_{XY} &= \{uv \mid u \in X \text{ and } v \in Y\}, \\
E_Y &= \{uv \mid u, v \in Y\}, \\
E_{ZXY} &= \{uv \mid u \in Z \text{ and } v \in X \cup Y\}.
\end{aligned}$$

Then

$$\begin{aligned} V(K_{n,n} - nK_2) &= X \cup Y \cup Z, \quad X \cap Y = X \cap Z = Y \cap Z = \emptyset. \\ E(K_{n,n} - nK_2) &= E_X \cup E_{XY} \cup E_Y \cup E_{ZY}. \end{aligned}$$

In drawings  $D_n$  with odd  $n \geq 5$ , vertices  $a_{2\lceil \frac{n-1}{4} \rceil - 1}, b_{2\lceil \frac{n-1}{4} \rceil}, \dots, a_{2\lceil \frac{n-1}{4} \rceil - 3}, b_{2\lceil \frac{n-1}{4} \rceil - 2}$  ( $a_0, b_1, \dots, a_{n-3}, b_{n-2}$ ) of  $X$  ( $Y$ ) are placed equidistantly on the perimeter of the top (bottom) disk, vertex  $a_{n-1}$  ( $b_{n-1}$ ) is placed on the center of the top (bottom) disk, edges of  $E_X$  ( $E_Y$ ) are drawn on the top (bottom) disk, and edges of  $E_{XY}$  are drawn by shortest helical curves on the cylinder. For  $0 \leq i \leq (n-3)/2$ , edges  $a_{n-1}b_{2i}$  ( $b_{n-1}a_{2i}$ ) are drawn on the top (bottom) disk, and edges  $a_{n-1}b_{2i+1}$  ( $b_{n-1}a_{2i+1}$ ) are drawn on the top (bottom) disk and cylinder. It follows  $\nu_{D_n}(E_X \cup E_Y \cup E_{XY}) = \nu(D_{n-1})$ ,  $\nu_{D_n}(E_{ZY}) = 0$  and

$$\begin{aligned} &\nu_{D_n}(E_X \cup E_Y \cup E_{XY}, E_{ZY}) \\ &= (n-1)(2\lceil \frac{n-1}{4} \rceil \lfloor \frac{n-3}{4} \rfloor + \lfloor \frac{n-1}{4} \rfloor + \sum_{d=0}^{\frac{n-5}{2}} d) \\ &= (n-1)(\lfloor \frac{n-1}{4} \rfloor (2\lceil \frac{n-3}{4} \rceil + 1) + \frac{n-5}{2} \frac{n-3}{2}) \\ &= (n-1)(\frac{(n-1)(n-3)}{8} + \frac{(n-3)(n-5)}{8}) \\ &= \frac{(n-1)(n-3)^2}{4}. \end{aligned}$$

By Lemma 2.2,

$$\begin{aligned} \nu(D_n) &= \nu_{D_n}(E_X \cup E_Y \cup E_{XY}) + \nu_{D_n}(E_X \cup E_Y \cup E_{XY}, E_{ZY}) + \nu_{D_n}(E_{ZY}) \\ &= \frac{(n-1)(n-3)^2(n-5)}{16} + \frac{(n-1)(n-3)^2}{4} \\ &= \frac{(n-1)^2(n-3)^2}{16} \\ &= \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor. \end{aligned}$$

So we have the following Lemma 2.3:

**Lemma 2.3.** For odd  $n \geq 5$ ,  $\nu(D_n) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ .

By Lemma 2.1-2.3, we have

**Theorem 2.1.** For  $n \geq 1$ ,  $cr(K_{n,n} - nK_2) \leq \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ .

Since  $K_n \times P_2 \cong K_{n,n} - nK_2$ , we have

**Corollary 2.1.** For  $n \geq 1$ ,  $cr(K_n \times P_2) \leq \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ .

### 3 Upper bound of $cr(K_n \times P_3)$

In this section, firstly we introduce some topological tools, and then we give a drawing for  $K_n \times P_3$  based on  $D_n$ .

We define a structure  $M_{l,r}^2$  in the real plane  $\mathbb{R}^2$ . For the vertical points  $(0, 0)$  and  $(0, 1)$ , let  $S_l = \{(l_i^0, l_i^1) : 0 \leq i \leq l-1\}$  ( $S_r = \{(r_i^0, r_i^1) : 0 \leq i \leq r-1\}$ ) be a set of non-vertical bunches of lines in the left (right) semi-plane, such that the point  $(0, j)$  belongs to  $l_i^j$  ( $r_i^j$ ) for  $0 \leq i \leq l-1$  ( $0 \leq i \leq r-1$ ) and  $j = 0, 1$ . In Figure 3.1 we show the drawings of  $M_{1,2}^2$  and  $M_{2,2}^2$  as examples.




Figure 3.1: Drawings of  $M_{1,2}^2$  and  $M_{2,2}^2$

By counting the crossings in  $M_{l,r}^2$ , we have

**Lemma 3.1.** *For positive integers  $l$  and  $r$ ,*

$$\nu_{M_{l,r}^2}(S_l) + \nu_{M_{l,r}^2}(S_r) = \binom{l}{2} + \binom{r}{2} = \frac{l(l-1)}{2} + \frac{r(r-1)}{2}.$$

Let

$$V(K_n \times P_3) = \{0i, 1i, 2i \mid 0 \leq i \leq n-1\},$$

$$E(K_n \times P_3) = \{(0i, 1j), (2i, 1j) \mid 0 \leq i \neq j \leq n-1\}.$$

We construct a drawing  $D'_n$  for  $K_n \times P_3$  based on  $D_n$  with  $M_{\frac{n-1}{2}, \frac{n-1}{2}}^2$  for odd  $n$  and  $M_{\frac{n}{2}, \frac{n-2}{2}}^2$  or  $M_{\frac{n-2}{2}, \frac{n}{2}}^2$  for even  $n$ . Vertex  $a_i$  ( $b_i$ ) ( $0 \leq i \leq n-1$ ) in its “small” neighborhood in the drawing  $D_n$  is replaced by two vertices  $0i$  and  $2i$  vertically (one vertex  $1i$ ). Now every drawn edge  $e$  in  $D_n$  which starts from  $a_i$  is replaced by a bunch of two edges described above, which is drawn along the original edge  $e$ . Drawings  $D'_n$  for  $n \leq 9$  are shown in Figure 3.2.

There are totally two types of crossings in  $D'_n$ .

**Type 1.** Crossings produced by the crossings in  $D_n$ .

Each crossing in  $D_n$  will be replaced by 4 crossings in  $D'_{n+1}$ , since we replace every edge with a bunch of two edges.

**Type 2.** Crossings produced by replacing a vertex of  $D_n$  by two vertices in  $D'_n$ .




Figure 3.2: Drawings  $D'_n$  for  $n \leq 9$

The total number of crossings of Type 2 are  $n \cdot \nu(M_{\frac{n-1}{2}, \frac{n-1}{2}}^2)$  for odd  $n$  and  $n \cdot \nu(M_{\frac{n}{2}, \frac{n-2}{2}}^2)$  for even  $n$ .

By Lemma 3.1, for odd  $n$ ,

$$\begin{aligned}
 \nu(D'_n) &= 4\nu(D_n) + n \cdot \nu(M_{\frac{n-1}{2}, \frac{n-1}{2}}^2) \\
 &= 4\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \left( \frac{\frac{n-1}{2} \frac{n-3}{2}}{2} + \frac{\frac{n-1}{2} \frac{n-3}{2}}{2} \right) \\
 &= 4\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor.
 \end{aligned}$$

For even  $n$ ,

$$\begin{aligned}
 \nu(D'_n) &= 4\nu(D_n) + n \cdot \nu(M_{\frac{n}{2}, \frac{n-2}{2}}^2) \\
 &= 4\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \left( \frac{\frac{n}{2} \frac{n-2}{2}}{2} + \frac{\frac{n-2}{2} \frac{n-4}{2}}{2} \right) \\
 &= 4\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor.
 \end{aligned}$$

Hence we have

**Theorem 3.1.** For  $n \geq 1$ ,  $cr(K_n \times P_3) \leq 4\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor$ .

## 4 Upper bound of $cr(K_n \times C_4)$

We define another structure  $M_{l,r}^4$  in the real plane  $\mathbb{R}^2$ . For the vertical points  $(0,0)$  and  $(0,1)$ , let  $S_l = \{(l_{i,0}^0, l_{i,1}^0, l_{i,0}^1, l_{i,1}^1) : 0 \leq i \leq l-1\}$  ( $S_r = \{(r_{i,0}^0, r_{i,1}^0, r_{i,0}^1, r_{i,1}^1) : 0 \leq i \leq r-1\}$ ) be a set of non-vertical bunches of lines in the left (right) semi-plane, such that the point  $(0,j)$  belongs to  $l_{i,k}^j$  ( $r_{i,k}^j$ ) for  $0 \leq i \leq l-1$  ( $0 \leq i \leq r-1$ ),  $j = 0, 1$  and  $k = 0, 1$ . In each bunch of lines of  $S_l$  ( $S_r$ ), there is exact one ‘self’ crossing crossed by  $l_{i,1}^0$  and  $l_{i,1}^1$  ( $r_{i,1}^0$  and  $r_{i,1}^1$ ). In Figure 4.1 we show the drawings of  $M_{1,2}^4$  and  $M_{2,2}^4$  as examples.




Figure 4.1: Drawings of  $M_{1,2}^4$  and  $M_{2,2}^4$

By counting the crossings in  $M_{l,r}^4$ , we have

**Lemma 4.1.** *For positive integers  $l$  and  $r$ ,*

$$\nu_{M_{l,r}^4}(S_l) + \nu_{M_{l,r}^4}(S_r) = \binom{2l}{2} + \binom{2r}{2} = l(2l-1) + r(2r-1).$$

Let

$$V(K_n \times C_4) = \{0i, 1i, 2i, 3i \mid 0 \leq i \leq n-1\},$$

$$E(K_n \times C_4) = \{(0i, 1j), (1i, 2j), (2i, 3j), (3i, 0j) \mid 0 \leq i \neq j \leq n-1\}.$$

We construct a drawing  $D_n''$  for  $K_n \times C_4$  based on  $D_n$  with  $M_{\frac{n-1}{2}, \frac{n-1}{2}}^4$  for odd  $n$  and  $M_{\frac{n}{2}, \frac{n-2}{2}}^4$  or  $M_{\frac{n-2}{2}, \frac{n}{2}}^4$  for even  $n$ . Vertex  $a_i$  ( $b_i$ ) ( $0 \leq i \leq n-1$ ) in its ‘small’ neighborhood in the drawing  $D_n$  is replaced by two vertices  $0i$  and  $2i$  ( $1i$  and  $3i$ ) vertically. Now every drawn edge  $e$  in  $D_n$  which starts from  $a_i$  ( $b_i$ ) is replaced by a bunch of four edges described above, which is drawn along the original edge  $e$ . For even  $n \geq 4$ , by drawing edge  $l_{i,1}^0$  (or  $l_{i,1}^1, r_{i,1}^0, r_{i,1}^1$ ) carefully,  $2n$  ‘self’ crossings can be reduced. Drawings  $D_n''$  for  $n \leq 9$  are shown in Figure 4.2.

Since  $K_1 \times C_4 \cong 4K_1$  and  $K_2 \times C_4 \cong 2C_4$ , we have  $cr(K_1 \times C_4) = cr(K_2 \times C_4) = 0$ . Now, we consider the cases of  $n \geq 3$ .




Figure 4.2: Drawings  $D''_n$  for  $n \leq 9$

There are totally two types of crossings in  $D''_n$ .

**Type 1.** Crossings produced by the crossings in  $D_n$ .

Each crossing in  $D_n$  will be replaced by 16 crossings in  $D''_{n+1}$ , since we replace every edge with a bunch of four edges.

**Type 2.** Crossings produced by replacing a vertex of  $D_n$  by two vertices in  $D''_n$ .

The total number of crossings of Type 2 are  $2n \cdot \nu(M_{\frac{n-1}{2}, \frac{n-1}{2}}^4) - n(n-1)$  for odd  $n$  and  $2n \cdot \nu(M_{\frac{n}{2}, \frac{n-2}{2}}^4) - n(n-1) - 2n$  for even  $n$ . Notice that we have to subtract  $|E(D_n)| = n(n-1)$  since the ‘self’ crossings of each bunch are counted twice (namely both of the endpoints).

By Lemma 4.1, for odd  $n \geq 3$ ,

$$\begin{aligned}\nu(D_n'') &= 16\nu(D_n) + 2n \cdot \nu(M_{\frac{n-1}{2}, \frac{n-1}{2}}^4) - n(n-1) \\ &= 16\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + 2n(n-1)(n-2) - n(n-1) \\ &= 16\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n(n-1)(2n-5).\end{aligned}$$

For even  $n \geq 4$ ,

$$\begin{aligned}\nu(D_n'') &= 16\nu(D_n) + 2n \cdot \nu(M_{\frac{n}{2}, \frac{n-2}{2}}^4) - n(n-1) - 2n \\ &= 16\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + 2n(n^2 - 3n + 3) - n(n-1) - 2n \\ &= 16\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n(n-1)(2n-5).\end{aligned}$$

Hence we have

**Theorem 4.1.** *For  $n \geq 3$ ,  $cr(K_n \times C_4) \leq 16\lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n(n-1)(2n-5)$ .*

## 5 Lower bounds of $cr(K_{n,n} - nK_2)$ , $cr(K_n \times P_2)$ , $cr(K_n \times P_3)$ and $cr(K_n \times C_4)$

We shall introduce the lower bound method proposed by Leighton [12]. Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be graphs. An embedding of  $G_1$  in  $G_2$  is a couple of mappings  $(\varphi, \kappa)$  satisfying

$\varphi : V_1 \rightarrow V_2$  is an injection

$\kappa : E_1 \rightarrow \{\text{set of all paths in } G_2\}$ ,

such that if  $uv \in E_1$  then  $\kappa(uv)$  is a path between  $\varphi(u)$  and  $\varphi(v)$ . For any  $e \in E_2$  define

$$cg_e(\varphi, \kappa) = |\{f \in E_1 : e \in \kappa(f)\}|$$

and

$$cg(\varphi, \kappa) = \max_{e \in E_2} \{cg_e(\varphi, \kappa)\}.$$

The value  $cg(\varphi, \kappa)$  is called congestion.

**Lemma 5.1.** [12] *Let  $(\varphi, \kappa)$  be an embedding of  $G_1$  in  $G_2$  with congestion  $cg(\varphi, \kappa)$ . Let  $\Delta(G_2)$  denote the maximal degree of  $G_2$ . Then*

$$cr(G_2) \geq \frac{cr(G_1)}{cg^2(\varphi, \kappa)} - \frac{|V_2|}{2} \Delta^2(G_2).$$

Let  $K_{m,n}^x$  be the complete bipartite multigraph of  $m + n$  vertices, in which every two vertices are joined by  $x$  parallel edges.

According to De Klerk [5] and Kainen [10], the following lemmas hold.

**Lemma 5.2.** [5]  $cr(K_{m,n}) \geq 0.8594 \lfloor \frac{m}{2} \rfloor \lfloor \frac{m-1}{2} \rfloor \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor$ .

**Lemma 5.3.** [10]  $cr(K_{m,n}^x) = x^2 cr(K_{m,n})$ .

Now we are in a position to show the lower bound of  $cr(K_{n,n} - nK_2)$ .

**Theorem 5.1.**  $cr(K_{n,n} - nK_2) \geq \frac{0.8594}{(1+\frac{3}{n-1})^2} \lfloor \frac{n}{2} \rfloor^2 \lfloor \frac{n-1}{2} \rfloor^2 - n(n-1)^2$ .

*Proof.* By Lemmas 5.1, 5.2 and 5.3, we only need to construct an embedding  $(\varphi, \kappa)$  of  $K_{n,n}^{(n-1)(n-2)}$  into  $K_{n,n} - nK_2$  with congestion  $cg(\varphi, \kappa) = (n-2)(n+2)$ .

Let  $\alpha_i^k \beta_i^k$  be the  $k$ -th  $(n-1, 2)$ -arrangement, where  $\alpha_i^k, \beta_i^k \in \{0, 1, 2, \dots, n-1\} - \{i\}$  and  $\alpha_i^k \neq \beta_i^k$ . Let

$$\begin{aligned} V(K_{n,n}^{(n-1)(n-2)}) &= \{u_i, v_i \mid 0 \leq i \leq n-1\}, \\ E(K_{n,n}^{(n-1)(n-2)}) &= \{(u_i, v_j)^k \mid 0 \leq i, j \leq n-1, 1 \leq k \leq (n-1)(n-2)\}. \end{aligned}$$

Let  $\varphi(u_i) = a_i$ ,  $\varphi(v_i) = b_i$ ,  $\kappa((u_i, v_j)^k) = P_{a_i b_{\alpha_i^k} a_{\beta_i^k} b_i}$  for  $0 \leq i \leq n-1$ , and  $\kappa((u_i, v_j)^k) = (a_i, b_j)$  for  $0 \leq i \neq j \leq n-1$ . Then  $cg_e(\varphi, \kappa) = (n-2)(n+2)$  for every  $e \in E(K_{n,n}^{(n-1)(n-2)})$ . This completes the proof of Theorem 5.1.  $\square$

Since  $K_n \times P_2 \cong K_{n,n} - nK_2$ , we have

**Corollary 5.1.**  $cr(K_n \times P_2) \geq \frac{0.8594}{(1+\frac{3}{n-1})^2} \lfloor \frac{n}{2} \rfloor^2 \lfloor \frac{n-1}{2} \rfloor^2 - n(n-1)^2$ .

Similar to the proof of Theorem 5.1, by constructing an embedding  $(\varphi, \kappa)$  of  $K_{2n,n}^{(n-1)(n-2)}$  into  $K_n \times P_3$  with congestion  $cg(\varphi, \kappa) = (n-2)(n+2)$ , we can get

**Theorem 5.2.**  $cr(K_n \times P_3) \geq \frac{0.8594}{(1+\frac{3}{n-1})^2} n \lfloor \frac{2n-1}{2} \rfloor \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor - \frac{3}{2} n(2n-2)^2$ .

And by constructing an embedding  $(\varphi, \kappa)$  of  $K_{2n,2n}^{(n-1)(n-2)}$  into  $K_n \times C_4$  with congestion  $cg(\varphi, \kappa) = (n-2)(n+2)$ , we can get

**Theorem 5.3.**  $cr(K_n \times P_4) \geq \frac{0.8594}{(1+\frac{3}{n-1})^2} n^2 \lfloor \frac{2n-1}{2} \rfloor^2 - 2n(2n-2)^2$ .

## 6 Conclusion

By Theorem 2.1, Theorem 3.1, Theorem 4.1 and Theorem 5.1-5.3, we have

$$\begin{aligned} \frac{0.8594}{(1 + \frac{3}{n-1})^2} \lfloor \frac{n}{2} \rfloor^2 \lfloor \frac{n-1}{2} \rfloor^2 - n(n-1)^2 &\leq cr(K_n \times P_2) = cr(K_{n,n} - nK_2) \leq \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor, \\ \frac{0.8594}{(1 + \frac{3}{n-1})^2} n \lfloor \frac{2n-1}{2} \rfloor \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor - \frac{3}{2} n(2n-2)^2 &\leq cr(K_n \times P_3) \leq 4 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor, \\ \frac{0.8594}{(1 + \frac{3}{n-1})^2} n^2 \lfloor \frac{2n-1}{2} \rfloor^2 - 2n(2n-2)^2 &\leq cr(K_n \times P_4) \leq 16 \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor + n(n-1)(2n-5) \text{ for } n \geq 3. \end{aligned}$$

From [6], we have Lemma 6.1

**Lemma 6.1.** *Let  $i$  be the least number of the edges of a graph  $G$  whose deletion from  $G$  results in a planar subgraph of  $G$ , then  $cr(G) \geq i$ .*

By Lemma 6.1, we have

**Lemma 6.2.**  $cr(K_{5,5} - 5K_2) = 4$ .

*Proof.* Let  $i$  be the least number of the edges of graph  $G = K_{5,5} - 5K_2$  whose deletion from  $G$  results in a planar subgraph  $G^*$  of  $G$ . Consider  $G^*$  has 10 vertices,  $20 - i$  edges. Let  $D^*$  be a planar drawing of  $G^*$  and  $f$  denote the number of the faces in  $D^*$ . Then according to the Euler Polyhedron Formula,

$$\begin{aligned} 10 - (20 - i) + f &= 2, \\ f &= 12 - i. \end{aligned}$$

Since the girth of  $G$  is 4, by counting the number of edges of each face in  $D^*$ , we have

$$\begin{aligned} (12 - i) \times 4 &\leq 2(20 - i), \\ i &\geq 4. \end{aligned}$$

By Lemma 6.1,  $cr(K_{5,5} - 5K_2) \geq 4$ . By Theorem 2.1,  $cr(K_{5,5} - 5K_2) \leq 4$ . So  $cr(K_{5,5} - 5K_2) = 4$ .  $\square$

Furthermore, we have the following conjecture.

**Conjecture 6.1.**  $cr(K_{n,n} - nK_2) = \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-1}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor \lfloor \frac{n-3}{2} \rfloor$ .

Conjecture 6.1 holds for  $n \leq 5$  by Lemma 2.1 and Lemma 6.2.

## References

- [1] J. Adamsson, Arrangements, circular arrangements and the crossing number of  $C_7 \square C_n$ , *J. Combin. Theory Ser. B* 90 (2004) 21-39.
- [2] D. Archdeacon, Topological graph: a survey, *Cong. numerantium* 115 (1996) 5-54.
- [3] L. Beineke, R. Welson, Selected topics in graph theory, *Academic Press* (1978) 68-72.
- [4] Z. Blažek, N. Koman, A Minimal problem concerning complete plane graphs, in M. Fiedler(Ed.) *Theory of Graphs and its Applications, Proceedings of the Symposium on Smolenice, 1963 Publ. House of the Czechoslovak Academy of Sciences, Prague* (1964) 118-117.
- [5] E. De Klerk, E. V. Pasechnik, A. Schrijver, Reduction of symmetric semidefinite programs using the regular representation, *Math Program, Ser B*, 109 (2007) 613-624.
- [6] S. Fiorini, On the crossing number of generalized Petersen graphs, *Ann, Discrete Math.* 30 (1986) 211-242.
- [7] M. R. Garey, D. S. Johnson, Crossing number is NP-complete, *SIAM J. Alg. Disc. Math.* 4 (1983) 312-316.
- [8] R. K. Guy, The decline and fall of Zarankiewicz's theorem, in F. Harary(Ed. ), *Proof Techniques in Graph Theory, Academic Press, New York, London* (1969) 63-69.
- [9] P. K. Jha, S. Devisetty, Orthogonal drawings and crossing numbers of the Kronecker product of two cycles *J. Parallel Distrib. Comput.* 72 (2012) 195-204.
- [10] P. C. Kainen, A lower bound for crossing numbers of graphs with applications to  $K_n$ ,  $K_{p,q}$ , and  $Q(d)$ . *J. Combin. Theory Ser. B* **12**, 287-298 (1972)
- [11] D. J. Kleitman, The crossing number of  $K_{5,n}$ , *J. Combinatorial Theory* 9 (1970) 315-323.
- [12] F. T. Leighton, New lower bound techniques for VLSI, *Math. Systems Theory* **17** (1984) 47-70.
- [13] D. R. Woodall, Cyclic-order graphs and Zarankiewicz's crossing-number conjecture, *J. Graph Theory* 17 (1993) 657-671.

- [14] K. Zarankiewicz, On a problem of P.Turán concerning graphs, *Fund Math* 41 (1954) 137-145.
- [15] W. Zheng, X. Lin, Y. Yang, C. Cui, On the crossing number of  $K_m \square P_n$ , *Graphs and Combinatorics* 23 (2007) 327-336.
- [16] W. Zheng, X. Lin, Y. Yang, The crossing number of  $K_{2,m} \square P_n$ , *Discrete Mathematics* 308 (2008) 6639-6644.
- [17] W. Zheng, X. Lin, Y. Yang, C. Deng, On the crossing number of  $K_m \square C_n$  and  $K_{m,l} \square P_n$ , *Discrete Applied Mathematics* 156 (2008) 1892-1907.